

PARTON DISTRIBUTIONS : A NEW GLOBAL ANALYSIS^a

A.D. MARTIN^a, R.G. ROBERTS^b, W.J. STIRLING^{a,c}, and R.S. THORNE^d

^a *Dept. of Physics, University of Durham, Durham, DH1 3LE*

^b *Rutherford Appleton Laboratory, Chilton, Didcot, Oxon, OX11 0QX*

^c *Dept. of Mathematical Sciences, University of Durham, Durham, DH1 3LE*

^d *Jesus College, University of Oxford, Oxford OX1 3DW*

We present a new analysis of parton distributions of the proton. This incorporates a wide range of new data, an improved treatment of heavy flavours and a re-examination of prompt photon production. The new set (MRST) shows systematic differences from previous sets of partons which can be identified with particular features of the new data and with improvements in the analysis. An ‘explanation’ of the behaviour seen on the Caldwell plot is offered.

1 New data

New experimental information used in the analysis includes the following.

- (i) New, more precise, measurements of the structure function F_2^{ep} for deep inelastic electron-proton scattering by the H1 and ZEUS collaborations at HERA ^{1,2}.
- (ii) A re-analysis of the CCFR neutrino data leading to a new set of $F_2^{\nu N}$ and $xF_3^{\nu N}$ structure function measurements ³.
- (iii) Measurements of the charm component of F_2^{ep} in electron-proton deep inelastic scattering at HERA ^{4,5}. These complement the existing EMC charm data ⁶.
- (iv) Very precise measurements of prompt photon production, from the E706 collaboration ⁷. These data motivate us to look again at our treatment of this reaction, and in particular of the WA70 prompt photon measurements ⁸. We emphasize that such prompt photon data are the main constraints on the gluon.
- (v) The E866 collaboration ⁹ have measured the asymmetry in Drell-Yan production in pp and pn collisions over an extended x range, $0.03 \lesssim x \lesssim 0.35$. The asymmetry data provide direct information on the x dependence of the difference, $\bar{u} - \bar{d}$, of the sea quark densities.

^aTalk by Dick Roberts at DIS98 workshop, Brussels, April 1998

- (vi) The new CDF¹¹ data on the asymmetry of the rapidity of the charged lepton from $W^\pm \rightarrow l^\pm \nu$ decays at the Tevatron $p\bar{p}$ collider. The new data extend to larger values of lepton rapidity.
- (vii) The final NMC analysis on $F_2^{\mu p}$, $F_2^{\mu d}$ and the ratio $F_2^{\mu d}/F_2^{\mu p}$ ¹².
- (viii) The data on Drell-Yan production obtained by the E772 collaboration¹³ cover a wider range of x_F than the E605 data¹⁴ which we have used to constrain the sea.

2 Improvements in the analysis

We parametrize the starting parton distributions at $Q^2 = Q_0^2 = 1 \text{ GeV}^2$ where the number of active flavours is $n_f = 3$. We work in the $\overline{\text{MS}}$ renormalization scheme and use the standard starting MRS parametric forms.

For the first time our treatment of the heavy flavour densities, charm and bottom, is on a firm theoretical footing. These densities are determined by the other parton distributions and no extra parameters are introduced apart from the heavy quark masses. At very low Q^2 the structure functions $F_2^H(x, Q^2)$, with $H = c, b$, are described by boson-gluon fusion and the heavy quark densities turn on at $Q^2 \simeq m_H^2$. The procedure is described in the talk by Robert Thorne¹⁵.

A special feature of our analysis is the particular attention to the uncertainties in the gluon distribution at large x . The main constraints in this region are data on prompt photon production in pp or pA collisions from the WA70⁸ and the E706⁷ experiments. We begin by taking a canonical value of $\langle k_T \rangle = 0.4 \text{ GeV}$ for the the WA70 data. We then explore a range of gluon distributions in which $\langle k_T \rangle$ goes from one extreme of $\langle k_T \rangle = 0$ to the other $\langle k_T \rangle = 0.64 \text{ GeV}$. We call the gluon distributions which correspond to $\langle k_T \rangle = 0, 0.4$ and 0.64 GeV the higher, central and lower (large x) gluons respectively — since a smaller gluon density is compensated by a larger $\langle k_T \rangle$. We denote the corresponding parton sets by MRST($g \uparrow$), MRST and MRST($g \downarrow$).

The optimum global MRST description has the QCD parameter $\Lambda_{\overline{\text{MS}}}(n_f = 4) = 300 \text{ MeV}$, which corresponds to $\alpha_S(M_Z^2) = 0.1175$.

The starting gluon and sea-quark distributions at small x , $xg \sim x^{-\lambda_g}$, $xS \sim x^{-\lambda_S}$ and we find negative values of λ_g , which imply a ‘valence’ type of behaviour for the gluon at low x for $Q^2 = 1 \text{ GeV}^2$. As Q^2 increases the valence-like character of the gluon rapidly disappears due to evolution being driven by the steeper sea, and by $Q^2 \simeq 2 \text{ GeV}^2$ the gluon is ‘flat’ in x , that is $\lambda_g = 0$.

Fig. 1 is our version of the ‘Caldwell plot’¹⁶ showing $\partial F_2/\partial \ln Q^2$ versus x . The curve shows the values of the slope $\partial F_2/\partial \ln Q^2$ versus x for the H1 data, compared with the slope found in the MRST fit (evaluated at the particular values of Q^2 appropriate to the experimental data). Thus the claim¹⁷ of a sudden change of dynamics at low x may be attributed simply to the rapidly varying nature of the gluon at very low Q^2 .




Figure 1: The description of the slopes $\partial F_2/\partial \ln Q^2$ versus x for the HERA data.

However, this ‘conventional’ explanation for the Caldwell plot may, in turn, be a signal for some breakdown of exact DGLAP around 1 GeV²; certainly we cannot move lower down in Q^2 as the gluon density (and hence F_L) starts to become negative at very low x .

3 Summary

This talk is an advertisement for the MRST parton sets. The full details of the analysis are available¹⁸. In addition to the sets MRST, MRST($g \uparrow$), MRST($g \downarrow$) which explore the likely allowed variation of the gluon we have sets MRST($\alpha_S \uparrow \uparrow$) and MRST($\alpha_S \downarrow \downarrow$) which explore a variation of α_S of 0.005 about the central value of 0.1175. These are all NLO sets evaluated in the $\overline{\text{MS}}$ scheme and are available from the Durham database (<http://durpdg.dur.ac.uk/HEPDATA>) together with a package which computes the relevant structure functions including F_L and F_2^{charm} . Furthermore, the corresponding sets evaluated in the DIS scheme can be likewise accessed.

References

1. H1 collaboration: S. Aid et al., Nucl. Phys. **B470** (1996) 3; C. Adloff et al., Nucl. Phys. **B497** (1997) 3.
2. ZEUS collaboration: M. Derrick et al., Zeit. Phys. **C69** (1996) 607; M. Derrick et al., Zeit. Phys. **C72** (1996) 399.
3. CCFR collaboration: W.G. Seligman et al., Phys. Rev. Lett. **79** (1997) 1213.
4. H1 collaboration: C. Adloff et al., Zeit. Phys. **C72** (1996) 593.
5. ZEUS collaboration: J. Breitweg et al., Phys. Lett. **B407** (1997) 402; Paper N-645 presented at International Europhysics Conference on High Energy Physics, HEP97, Jerusalem 1997.
6. J.J. Aubert et al., Nucl. Phys. **B213** (1983) 31.
7. E706 collaboration: L. Apanasevich et al., [hep-ex/9711017](#).
8. WA70 collaboration: M. Bonesini et al., Z. Phys. **C38** (1988) 371.
9. E866 collaboration: E.A. Hawker et al., Phys. Rev. Lett. **80** (1998) 3715.
10. NA51 collaboration: A. Baldit et al., Phys. Lett. **B332** (1994) 244.
11. CDF collaboration: presented by A. Bodek at the Cracow International Symposium on Radiative Corrections (CRAD96), Cracow, Poland, August 1–5, 1996, Acta Phys. Polon. **B28** (1997) 477.
12. NMCollaboration: M. Arneodo et al., Nucl. Phys. **B483** (1997) 3.
13. E772 collaboration: P.L. McGaughey et al., Phys. Rev. **D50** (1994) 3038.
14. E605 collaboration: G. Moreno et al., Phys. Rev. **D43** (1991) 2815.
15. R.S. Thorne, talk at this WG1 session.
R.S. Thorne and R.G. Roberts, [hep-ph/9709442](#) to be published in Phys. Rev. **D 57** vol 11.
R.S. Thorne and R.G. Roberts, Phys. Lett. **B421** (1998) 303.
16. A. Caldwell, DESY Theory Workshop, Sept. 1997.
17. A.H. Mueller, Opening plenary talk at this workshop.
18. A.D. Martin, R.G. Roberts, W.J. Stirling and R.S. Thorne, [hep-ph/9803445](#), to be published in Eur. Jour. Phys. A.